

BCFW recursion for amplitudes with off-shell gluons

Andreas van Hameren

The Henryk Niewodniczański Institute of Nuclear Physics
Polish Academy of Sciences

presented at

LoopFest XIII, City College of Technology, New York, 19-06-2014



This work was partially supported by
NCBiR grant LIDER/02/35/L-2/10/NCBiR/2011




Forward–central dijet decorrelations

van Hameren, Kotko,
Kutak, Sapeta 2014


2 jets

one initial state off–shell




3 jets

one initial state off–shell



both initial states on–shell



Hybrid “ k_T -collinear” factorization formula:

$$d\sigma_{AB \rightarrow X} = \int \frac{d^2 k_T}{\pi} \int \frac{dx_A}{x_A} \int dx_B \sum_b \mathcal{F}_{g^*/A}(x_A, k_T, \mu) f_{b/B}(x_B, \mu) d\hat{\sigma}_{g^* b \rightarrow X}(x_A, x_B, k_T, \mu)$$

The 2-jet calculation cannot be performed at LO within collinear factorization only and without a parton shower.

High-energy factorization

Collins, Ellis 1991

Catani, Ciafaloni, Hautmann 1991

$$\sigma_{h_1, h_2 \rightarrow QQ} = \int d^2 k_{1\perp} \frac{dx_1}{x_1} \mathcal{F}(x_1, k_{1\perp}) d^2 k_{2\perp} \frac{dx_2}{x_2} \mathcal{F}(x_2, k_{2\perp}) \hat{\sigma}_{gg} \left(\frac{m^2}{x_1 x_2 s}, \frac{k_{1\perp}}{m}, \frac{k_{2\perp}}{m} \right)$$

- to be applied in the 3-scale regime $s \gg m^2 \gg \Lambda_{\text{QCD}}^2$
- reduces to collinear factorization for $s \gg m^2 \gg k_\perp^2$,
but holds also for $s \gg m^2 \sim k_\perp^2$
- *unintegrated pdf* \mathcal{F} may satisfy BFKL-eqn, CCFM-eqn, BK-eqn, KGBJS-eqn, ...
- typically associated with small- x physics
- relevant for forward physics, saturation physics, heavy-ion physics...
- k_\perp gives a handle on the size of the proton
- allows for higher-order kinematical effects at leading order

Scattering amplitudes with off-shell legs

How to define and calculate scattering amplitudes with off-shell legs? How to ensure *gauge invariance*?

- Lipatov's effective action, in terms of two extra fields, so-called *reggeons*.

Lipatov 1995

Antonov, Lipatov, Kuraev, Cherednikov 2005


- Determine extra terms to be added to the amplitude using Slavnov-Taylor identities.

AvH, Kotko, Kutak 2012


- Embed the gluon scattering process into a quark scattering process, where the auxiliary quarks satisfy eikonal Feynman rules.

AvH, Kotko, Kutak 2013

$$k_1 = x_1 \ell_1 + k_{1\perp}$$



$$k_2 = x_2 \ell_2 + k_{2\perp}$$



Off-shell gluons from Wilson lines

Kotko 2014

A Wilson line along path C , defined as

$$[x, y]_C = \mathcal{P} \exp \left\{ ig \int_C dz_\mu A_b^\mu(z) T^b \right\},$$

transforms under local gauge transformations as $[x, y]_C \mapsto U(x)[x, y]_C U^\dagger(y)$.
Use an infinite Wilson line with “direction” p^μ

$$[y]_p = \mathcal{P} \exp \left\{ ig \int_{-\infty}^{\infty} ds p \cdot A_b(y + sp) T^b \right\}$$

to define the operator


$$\mathcal{R}^a(p, k) = \int d^4y e^{iy \cdot k} \text{Tr} \left\{ \frac{1}{\pi g} T^a [y]_p \right\}.$$

Amplitudes with n on-shell gluons and m off-shell gluons defined by

$$\begin{aligned} & \langle k_1, k_2, \dots, k_n | \mathcal{R}^{a_{n+1}}(p_{n+1}, k_{n+1}) \mathcal{R}^{a_{n+2}}(p_{n+2}, k_{n+2}) \cdots \mathcal{R}^{a_{n+m}}(p_{n+m}, k_{n+m}) | 0 \rangle \\ &= \delta(p_{n+1} \cdot k_{n+1}) \delta(p_{n+2} \cdot k_{n+2}) \cdots \delta(p_{n+m} \cdot k_{n+m}) \delta^4(k_1 + k_2 + \cdots + k_{n+m}) \\ & \times \mathcal{A}(k_1, k_2, \dots, k_{n+m}; p_{n+1}, p_{n+2}, \dots, p_{n+m}) \end{aligned}$$

Feynman rules

Planar graphs for the process $\emptyset \rightarrow g^*g^*g$:



The Feynman rules in the Feynman gauge:

$$\mu \text{ --- } \nu = \frac{-\eta^{\mu\nu}}{K^2} \quad \text{---} = \frac{1}{2p \cdot K} \quad \overline{\text{---}}_{\mu} = \sqrt{2} p^{\mu}$$

$$\begin{array}{c}
 \text{---} \\
 | \\
 1 \quad 2 \quad 3
 \end{array}
 = \frac{1}{\sqrt{2}} [(K_1 - K_2)^{\mu_3} \eta^{\mu_1 \mu_2} + (K_2 - K_3)^{\mu_1} \eta^{\mu_2 \mu_3} + (K_3 - K_1)^{\mu_2} \eta^{\mu_3 \mu_1}]$$

$$\begin{array}{c}
 \text{---} \\
 | \\
 1 \quad 2 \quad 3 \\
 \diagup \quad \diagdown \\
 | \\
 4
 \end{array}
 = \frac{-1}{2} [2 \eta^{\mu_1 \mu_3} \eta^{\mu_2 \mu_4} - \eta^{\mu_1 \mu_2} \eta^{\mu_3 \mu_4} - \eta^{\mu_1 \mu_4} \eta^{\mu_2 \mu_3}]$$

where p^{μ} is the direction associated with the eikonal line.

BCFW recursion

Britto, Cachazo, Feng 2004
 Britto, Cachazo, Feng, Witten 2005

For a rational function f of a complex variable z which vanishes at infinity, we have

$$\lim_{z \rightarrow \infty} f(z) = 0 \quad \Rightarrow \quad \oint \frac{dz}{2\pi i} \frac{f(z)}{z} = 0 ,$$

where the integration contour expands to infinity and necessarily encloses all poles of f . This directly leads to the relation

$$f(0) = \sum_i \frac{\lim_{z \rightarrow z_i} f(z)(z - z_i)}{-z_i} ,$$

where the sum is over all poles of f , and z_i is the position of pole number i . For color-ordered tree-level multi-gluon amplitudes, this can be translated to

$$\mathcal{A}(1^+, 2, \dots, n-1, n^-) = \sum_{i=2}^{n-1} \sum_{h=+,-} \mathcal{A}(\hat{1}^+, 2, \dots, i, -\hat{K}_{1,i}^h) \frac{1}{\hat{K}_{1,i}^2} \mathcal{A}(\hat{K}_{1,i}^h, i+1, \dots, n-1, \hat{n}^-)$$

where the lower-point on-shell amplitudes have “shifted” momenta.

BCFW for off-shell gluons

n -gluon amplitude is a function of n momenta k_1, k_2, \dots, k_n and n directions p_1, p_2, \dots, p_n , satisfying the conditions

$$\begin{aligned} k_1^\mu + k_2^\mu + \cdots + k_n^\mu &= 0 && \text{momentum conservation} \\ p_1^2 = p_2^2 = \cdots = p_n^2 &= 0 && \text{light-likeness} \\ p_1 \cdot k_1 = p_2 \cdot k_2 = \cdots = p_n \cdot k_n &= 0 && \text{eikonal condition} \end{aligned}$$

With the help of an auxiliary four-vector q^μ with $q^2 = 0$, we define

$$k_T^\mu(q) = k^\mu - x(q)p^\mu \quad \text{with} \quad x(q) \equiv \frac{q \cdot k}{q \cdot p}$$

Construct k_T^μ explicitly in terms of p^μ and q^μ :

$$k_T^\mu(q) = -\frac{\kappa}{2} \frac{\langle p | \gamma^\mu | q \rangle}{[pq]} - \frac{\kappa^*}{2} \frac{\langle q | \gamma^\mu | p \rangle}{\langle qp \rangle} \quad \text{with} \quad \kappa = \frac{\langle q | \kappa | p \rangle}{\langle qp \rangle}, \quad \kappa^* = \frac{\langle p | \kappa | q \rangle}{[pq]}$$

$k^2 = -\kappa\kappa^*$ is independent of q^μ , but also individually

κ and κ^* are independent of q^μ .

BCFW for off-shell gluons

n -gluon amplitude is a function of n momenta k_1, k_2, \dots, k_n and n directions p_1, p_2, \dots, p_n , satisfying the conditions

$$\begin{aligned} k_1^\mu + k_2^\mu + \cdots + k_n^\mu &= 0 && \text{momentum conservation} \\ p_1^2 = p_2^2 = \cdots = p_n^2 &= 0 && \text{light-likeness} \\ p_1 \cdot k_1 = p_2 \cdot k_2 = \cdots = p_n \cdot k_n &= 0 && \text{eikonal condition} \end{aligned}$$

With the help of an auxiliary four-vector q^μ with $q^2 = 0$, we define

$$\frac{\langle q|\kappa|p\rangle}{\langle qp\rangle} = \frac{\langle q|\kappa|p\rangle\langle pr\rangle}{\langle qp\rangle\langle pr\rangle} = \frac{\langle q|\kappa p|r\rangle}{\langle qp\rangle\langle pr\rangle} = \frac{\langle q|2k\cdot p - p\kappa|r\rangle}{\langle qp\rangle\langle pr\rangle} = -\frac{\langle qp\rangle[p|\kappa|r\rangle}{\langle qp\rangle\langle pr\rangle} = \frac{\langle r|\kappa|p\rangle}{\langle rp\rangle}$$

Construct k_T^μ explicitly in terms of p^μ and q^μ :

$$k_T^\mu(q) = -\frac{\kappa}{2} \frac{\langle p|\gamma^\mu|q\rangle}{[pq]} - \frac{\kappa^*}{2} \frac{\langle q|\gamma^\mu|p\rangle}{\langle qp\rangle} \quad \text{with} \quad \kappa = \frac{\langle q|\kappa|p\rangle}{\langle qp\rangle}, \quad \kappa^* = \frac{\langle p|\kappa|q\rangle}{[pq]}$$

$k^2 = -\kappa\kappa^*$ is independent of q^μ , but also individually

κ and κ^* are independent of q^μ .

BCFW: shifted momenta

$$f(0) = \sum_i \frac{\lim_{z \rightarrow z_i} f(z)(z - z_i)}{-z_i}$$

We choose two external gluons i and j , and use the direction of one as the auxiliary momentum to define the transverse momentum of the other, and vice versa, so

$$k_i^\mu = x_i(p_j)p_i^\mu - \frac{\kappa_i}{2} \frac{\langle i|\gamma^\mu|j\rangle}{[ij]} - \frac{\kappa_i^*}{2} \frac{\langle j|\gamma^\mu|i\rangle}{\langle ji\rangle}, \quad k_j^\mu = x_j(p_i)p_j^\mu - \frac{\kappa_j}{2} \frac{\langle j|\gamma^\mu|i\rangle}{[ji]} - \frac{\kappa_j^*}{2} \frac{\langle i|\gamma^\mu|j\rangle}{\langle ij\rangle}$$

We adopt the notation $|i\rangle \equiv |p_i\rangle$, $|j\rangle \equiv |p_j\rangle$ etc.. Choose the shift vector

$$e^\mu \equiv \frac{1}{2}\langle i|\gamma^\mu|j\rangle \quad \Rightarrow \quad p_i \cdot e = p_j \cdot e = e \cdot e = 0$$

Using this vector, we then define the shifted momenta

$$\begin{aligned} \hat{k}_i^\mu(z) &\equiv k_i^\mu + ze^\mu = x_i(p_j)p_i^\mu - \frac{\kappa_i - [ij]z}{2} \frac{\langle i|\gamma^\mu|j\rangle}{[ij]} - \frac{\kappa_i^*}{2} \frac{\langle j|\gamma^\mu|i\rangle}{\langle ji\rangle} \\ \hat{k}_j^\mu(z) &\equiv k_j^\mu - ze^\mu = x_j(p_i)p_j^\mu - \frac{\kappa_j}{2} \frac{\langle j|\gamma^\mu|i\rangle}{[ji]} - \frac{\kappa_j^* + \langle ij\rangle z}{2} \frac{\langle i|\gamma^\mu|j\rangle}{\langle ij\rangle} \end{aligned}$$

Total momentum is conserved, and also

$$p_i \cdot \hat{k}_i = 0 \quad , \quad p_j \cdot \hat{k}_j = 0$$

BCFW recursion

$$f(0) = \sum_i \frac{\lim_{z \rightarrow z_i} f(z)(z - z_i)}{-z_i}$$

The BCFW recursion formula becomes

$$\begin{array}{c} \cdot \cdot \cdot \\ \text{2} = \textcircled{=} \text{n-1} \\ | \quad | \\ 1 \quad n \end{array} = \sum_{i=2}^{n-2} \sum_{h=+,-} A_{i,h} + \sum_{i=2}^{n-1} B_i + C + D,$$

where

$$A_{i,h} = \begin{array}{c} i \\ \textcircled{=} \\ h \\ | \\ \hat{1} \end{array} \frac{1}{\kappa_{1,i}^2} \begin{array}{c} i+1 \\ \textcircled{=} \\ -h \\ | \\ \hat{n} \end{array}$$

$$B_i = \begin{array}{c} i-1 \\ \textcircled{=} \\ i \\ | \\ \hat{1} \end{array} \frac{1}{2p_i \cdot \kappa_{i,n}} \begin{array}{c} i \\ \textcircled{=} \\ i+1 \\ | \\ \hat{n} \end{array}$$

$$C = \frac{1}{\kappa_1} \begin{array}{c} \cdot \cdot \cdot \\ \text{2} = \textcircled{=} \text{n-1} \\ | \quad | \\ \hat{1} \quad \hat{n} \end{array}$$

$$D = \frac{1}{\kappa_n^*} \begin{array}{c} \cdot \cdot \cdot \\ \text{2} = \textcircled{=} \text{n-1} \\ | \quad | \\ \hat{1} \quad \hat{n} \end{array}$$

The hatted numbers label the shifted external gluons.

On-shell limit

For each off-shell gluon j , we can identify the following terms in the amplitude

$$\mathcal{A}(k_j) = \frac{1}{\kappa_j^*} U(k_j) + \frac{1}{\kappa_j} V(k_j) + W(k_j)$$

The actual amplitude needs a factor proportional to $\sqrt{-k_j^2}$, we choose κ_j^* :

$$\kappa_j^* \mathcal{A}(k_j) = U(k_j) + \frac{\kappa_j^*}{\kappa_j} V(k_j) + \kappa_j^* W(k_j)$$

The ratio κ_j^*/κ_j does not vanish in the on-shell limit, and an angle dependence remains.

$$|\kappa_j^* \mathcal{A}(k_j)|^2 \xrightarrow{k_j^2 \rightarrow 0} |U(p_j)|^2 + |V(p_j)|^2 + e^{2i\varphi_j} U(p_j)V(p_j)^* + e^{-2i\varphi_j} U(p_j)^*V(p_j)$$

Interference terms vanish upon integration over φ .

- the $-$ helicity can be associated with U , or $1/\kappa_j^*$
- the $+$ helicity can be associated with V , or $1/\kappa_j$

3-gluon amplitudes

$\emptyset \rightarrow ggg$, at least two shifted momenta

$$\mathcal{A}(1^+, 2^-, 3^-) = \frac{\langle 23 \rangle^3}{\langle 31 \rangle \langle 12 \rangle} \quad , \quad \mathcal{A}(1^-, 2^+, 3^+) = \frac{[32]^3}{[21][13]}$$

$\emptyset \rightarrow g^* gg$

$$\mathcal{A}(1^*, 2^+, 3^-) = \frac{1}{\kappa_1^*} \frac{\langle 31 \rangle^3}{\langle 12 \rangle \langle 23 \rangle} = \frac{1}{\kappa_1} \frac{[21]^3}{[13][32]}$$

$\emptyset \rightarrow gg^*g^*$

$$\mathcal{A}(1^+, 2^*, 3^*) = \frac{1}{\kappa_2^* \kappa_3^*} \frac{\langle 23 \rangle^3}{\langle 31 \rangle \langle 12 \rangle} \quad , \quad \mathcal{A}(1^-, 2^*, 3^*) = \frac{1}{\kappa_2 \kappa_3} \frac{[32]^3}{[21][13]}$$

$\emptyset \rightarrow g^*g^*g^*$

$$\mathcal{A}(1^*, 2^*, 3^*) = \frac{\langle 12 \rangle^3 [32]^3}{\kappa_3 \kappa_1^* \langle 1 | \kappa_3 | 2 \rangle \langle 2 | \kappa_1 | 3 \rangle \langle 2 | \kappa_1 | 2 \rangle} + (231) + (312) \, ,$$

where the second and third term are obtained by applying the cyclic permutations on the arguments and indices of the first term.

4-gluon amplitudes, 2 off-shell

$$\mathcal{A}(1^*, 2^+, 3^+, 4^*) = \frac{1}{\kappa_4^* \kappa_1^*} \frac{\langle 41 \rangle^3}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle} \quad , \quad \mathcal{A}(1^*, 2^+, 3^*, 4^+) = \frac{1}{\kappa_1^* \kappa_3^*} \frac{\langle 13 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle}$$

$$\begin{aligned} \mathcal{A}(1^*, 2^+, 3^-, 4^*) &= \frac{1}{\kappa_1^* \kappa_4} \frac{-\langle 1 | \not{p}_3 + \not{k}_4 | 4 \rangle^4}{\langle 2 | \not{k}_1 | 4 \rangle \langle 1 | \not{k}_4 | 3 \rangle \langle 12 \rangle [43] (\not{p}_3 + \not{k}_4)^2} \\ &+ \frac{1}{\kappa_1} \frac{\langle 34 \rangle^3 [14]^3}{\langle 4 | \not{k}_4 + \not{k}_1 | 1 \rangle \langle 2 | \not{k}_1 | 4 \rangle \langle 4 | \not{k}_1 | 4 \rangle \langle 23 \rangle} + \frac{1}{\kappa_4^*} \frac{[21]^3 \langle 14 \rangle^3}{\langle 4 | \not{k}_4 + \not{k}_1 | 1 \rangle \langle 1 | \not{k}_4 | 3 \rangle \langle 1 | \not{k}_4 | 1 \rangle [32]} \end{aligned}$$

$$\begin{aligned} \mathcal{A}(1^*, 2^-, 3^*, 4^+) &= \frac{\langle 13 \rangle^3 [13]^3}{\langle 34 \rangle \langle 41 \rangle \langle 1 | \not{k}_3 + \not{p}_4 | 3 \rangle \langle 3 | \not{k}_1 + \not{p}_4 | 1 \rangle [32] [21]} \\ &+ \frac{1}{\kappa_1^* \kappa_3} \frac{\langle 12 \rangle^3 [43]^3}{\langle 2 | \not{k}_3 | 4 \rangle \langle 1 | \not{k}_3 + \not{p}_4 | 3 \rangle (\not{k}_3 + \not{p}_4)^2} + \frac{1}{\kappa_1 \kappa_3^*} \frac{\langle 23 \rangle^3 [14]^3}{\langle 2 | \not{k}_1 | 4 \rangle \langle 3 | \not{k}_1 + \not{p}_4 | 1 \rangle (\not{k}_1 + \not{p}_4)^2} \end{aligned}$$

MHV amplitudes

$$\mathcal{A}(1^-, i^-, (\text{the rest})^+) = \frac{\langle p_1 p_i \rangle^4}{\langle p_1 p_2 \rangle \langle p_2 p_3 \rangle \cdots \langle p_{n-2} p_{n-1} \rangle \langle p_{n-1} p_n \rangle \langle p_n p_1 \rangle}$$

$$\mathcal{A}(1^+, i^+, (\text{the rest})^-) = \frac{[p_i p_1]^4}{[p_1 p_n] [p_n p_{n-1}] [p_{n-1} p_{n-2}] \cdots [p_3 p_2] [p_2 p_1]}$$

$$\mathcal{A}(1^*, i^-, (\text{the rest})^+) = \frac{1}{\kappa_1^*} \frac{\langle p_1 p_i \rangle^4}{\langle p_1 p_2 \rangle \langle p_2 p_3 \rangle \cdots \langle p_{n-2} p_{n-1} \rangle \langle p_{n-1} p_n \rangle \langle p_n p_1 \rangle}$$

$$\mathcal{A}(1^*, i^+, (\text{the rest})^-) = \frac{1}{\kappa_1} \frac{[p_i p_1]^4}{[p_1 p_n] [p_n p_{n-1}] [p_{n-1} p_{n-2}] \cdots [p_3 p_2] [p_2 p_1]}$$

$$\mathcal{A}(1^*, i^*, (\text{the rest})^+) = \frac{1}{\kappa_1^* \kappa_i^*} \frac{\langle p_1 p_i \rangle^4}{\langle p_1 p_2 \rangle \langle p_2 p_3 \rangle \cdots \langle p_{n-2} p_{n-1} \rangle \langle p_{n-1} p_n \rangle \langle p_n p_1 \rangle}$$

$$\mathcal{A}(1^*, i^*, (\text{the rest})^-) = \frac{1}{\kappa_1 \kappa_i} \frac{[p_i p_1]^4}{[p_1 p_n] [p_n p_{n-1}] [p_{n-1} p_{n-2}] \cdots [p_3 p_2] [p_2 p_1]}$$